# Inference of dynamic weighted cumulative residual entropy for Burr XII distribution based on progressive censoring

Amal S. Hassan<sup>1</sup>, E. A. Elsherpieny<sup>2</sup>, Wesal E. Aghel<sup>3</sup>

### Abstract

The dynamic weighted cumulative residual (DWCR) entropy is regarded as an additional measure of uncertainty related to the residual lifetime function in several disciplines, including survival analysis and reliability. This article presents the DWCR formula based on Havarda and Charvat. This measurement is called the DWCR Havarda and Charvat entropy (DWCRHCE). This work uses progressive Type II censoring to investigate the implications of DWCR Tsallis entropy (DWCRTE), DWCR Rényi entropy (DWCRRE), and DWCRHCE for the Burr XII distribution. Both classical and Bayesian methods are used to derive the estimators of these entropy metrics. Assuming independent gamma priors, we get the Bayes estimator of the suggested measures. Due to the lack of explicit forms, the Metropolis-Hastings approach was offered to determine the Bayes estimates for symmetric and asymmetric loss functions. To determine the efficacy of the suggested estimating techniques, several simulations were run for different censoring schemes. The simulation analysis leads us to the conclusion that, under a precautionary loss function followed by a linear exponential loss function, the Bayesian estimates of DWCRTE are generally more effective than the DWCRHCE or DWCRRE. Compared to maximum likelihood estimates, Bayesian estimates are preferred for different metrics. After that, a detailed explanation of the process is provided by looking at real data. The analysis of real-world data, specifically the Shasta reservoir water capacity data, aligns with the findings from simulated data. Notably, these findings have crucial implications for effective water resource management decisions.

Key words: Burr XII distribution, dynamic weighted cumulative residual entropy, Bayesian estimators, precautionary loss function.

Mathematical Subject Classification: 62F10.

© Amal S. Hassan, E. A. Elsherpieny, Wesal. E. Aghel. Article available under the CC BY-SA 4.0 licence 💽 🕐 🎯



<sup>&</sup>lt;sup>1</sup> Cairo University, Faculty of Graduate Studies for Statistical Research, Egypt. E-mail: amal52\_soliman@cu.edu.eg. ORCID: https://orcid.org/0000-0003-4442-8458.

<sup>&</sup>lt;sup>2</sup> Cairo University, Faculty of Graduate Studies for Statistical Research, Egypt.

E-mail: Elsayed.Elsherpieny@cu.edu.eg. ORCID: https://orcid.org/0000-0002-2635-8081.

<sup>&</sup>lt;sup>3</sup> Corresponding Author. Zawia University Faculty Sciences, Libya. E-mail: Wesalagil@yahoo.com, 12422022546163@pg.cu.edu.eg. ORCID: https://orcid.org/0009-0000-7246-7660.

## 1. Introduction

### 1.1. Progressive Type II censoring

Censorship is common in many fields, including pharmacology, social economics, and engineering, particularly in reliability and survival analysis (Wang & Gui, 2021). Due to time and cost constraints, it is difficult to completely observe the sample data in actual production. Even though they have been thoroughly examined, conventional censoring systems such as Type-I, Type-II, and hybrid schemes are inflexible, in that units cannot be removed arbitrarily. Cohen (1963) proposed the progressive censoring scheme (PCS) in order to overcome this restriction. In the PCS, units are removed from the experiment at different time points, with the number of units eliminated at each time point specified in advance.

The progressive type-II censoring (PT-IIC) method is one of the most popular censoring schemes. Here, we provide its description. Assume that *m* failures will be noticed when *n* identical units are put through a test. At the moment of the initial failure  $(y_{(1)})$ , the number  $r_1$  of the surviving units (n - 1) is randomly selected and removed from the experiment. At the second failure  $(y_{(2)})$ , the number  $r_2$  of the surviving units  $(n - r_1-2)$ is randomly selected and removed from the experiment, and so on, until the *m*<sup>th</sup> failure  $(y_{(m)})$  occurs, at which point all the remaining  $n - m - r_1 - r_2 - \cdots - r_{m-1}$  units are removed. Thus, the PCS includes *m* observed samples of failure  $y_{(1)} \leq y_{(2)} \leq \ldots \leq y_{(m)}$ , and survival items  $\mathbf{r} = (r_1, r_2, \cdots, r_m)$  such that  $n = m + r_1 + \cdots + r_m$ . Note that in the PCS,  $\mathbf{r} = (r_1, r_2, \cdots, r_m)$  is prefixed. Balakrishnan and Aggrawala (2000) offered historical context and a comprehensive overview of PCS. Notably, the following special cases can be noticed:

- Classical Type-II censoring: This occurs when  $r_1 = r_2 = \cdots = r_{m-1} = 0$  and  $r_m = n m$ .
- Complete sample: This occurs when m = n and  $r_i = 0$ ,  $i = 1, 2, \dots, n$ .

#### 1.2. Entropy measures

Shannon (1948) proposed the concept of entropy as a metric for quantifying uncertainty. Nowadays, the fields of economics, physics, telecommunications, communication theory, and reliability have given this criterion significant consideration. One parameter generalization of the Shannon entropy that may be applied as a randomness metric is the Rényi entropy. Numerous disciplines, including biology, genetics, electrical engineering, computer science, economics, chemistry and physics use Rényi entropy in their work. The entropy function as an extension of the Shannon entropy was first presented by Rényi (1961), followed by Havrda and Charvat (1967). The Rényi and Havrda and Charvat entropy of order c are defined respectively by the following expressions:

$$R(c) = \frac{1}{1-c} \log \left( \int_{-\infty}^{\infty} (f(y))^c \, dy \right); c > 0, c \neq 1,$$
  
$$H(c) = \left( 2^{1-c} - 1 \right)^{-1} \left( \int_{-\infty}^{\infty} (f(y))^c - 1 \right); \quad c > 0, c \neq 1.$$

Tsallis (1988) also presented the ideas of Tsallis entropy as a measure of quantifying randomness. The entropy can be used to determine the level of uncertainty associated with a random observation. Tsallis later utilized its distinctive characteristics and situated it within a physical framework. This metric is designated for any continuous random variable of order c, where and it is defined as follows:

$$T(c) = \frac{1}{(c-1)} \left( 1 - \int_{-\infty}^{\infty} f^c(y) dy \right); \quad c > 0, c \neq 1$$

Recently, entropy measure estimation utilizing different statistical distributions and sampling strategies has been studied by several writers (see, for example, Baratpour et al., 2007; Abo-Eleneen, 2011; Cho et al., 2015; Lee, 2017; Hassan and Zaky, 2019, 2021; Helmy et al., 2021; Hassan et al., 2022; Helmy et al., 2023; and Hassan at al., 2024b).

Different measures of uncertainty for probability distributions have attracted many writers, especially in works related to reliability analysis and survival. In light of Rényi's entropy's utility, Sunoj and Linu (2012) presented cumulative residual Rényi entropy (CRRE) of order *c* as below:

$$\bar{R}(c) = \frac{1}{(1-c)} \log\left(\int_{t}^{\infty} (\bar{F}(y))^{c} dy\right); c > 0, c \neq 1,$$

where  $\overline{F}(y) = 1 - F(y)$  is the survival function (SF). Additionally, Sunoj and Linu (2012) examined the primary characteristics of the dynamic version of the CRRE (DCRRE), which is extended to the residual lifetime  $Y_t = (Y - t | Y > t)$  based on survival function, rather than using probability density function (PDF), and found it to be beneficial for reliability modeling. The DCRRE of order *c* is given by:

$$\overline{R}^*(c) = (1-c)^{-1} \log\left(\frac{1}{(\overline{F}(t))^c} \int_t^\infty (\overline{F}(y))^c dy\right); \quad c > 0, c \neq 1.$$
(1)

Sati and Gupta (2015) established the DCR Tsallis entropy (DCRTE) as follows:

$$\overline{T}^*(c) = (c-1)^{-1} \left( 1 - (\overline{F}(t))^{-c} \int_t^\infty (\overline{F}(y))^c dy \right); \quad c > 0, c \neq 1,$$

In the literature, some statistical inferences based on the above dynamic entropy measures and their related works have been considered by several authors. For the Pareto distribution, Bayesian estimates (BEs) of the DCR entropy under various sampling conditions have been examined by several researchers (see Renjini et al., 2016a, 2016b, and 2018; and Ahmadini et al., 2020). The Lindley distribution's BE of DCRRE was examined by Almarashi et al. (2021). Al-Babtain et al. (2021) supplied the maximum likelihood estimates (MLEs) and BEs of the DCRRE for the Lomax distribution. Mohamed (2022) used generalized order statistics to study DCRTE and cumulative residual Tsallis entropy. The BEs of the DCRTE for the moment exponential distribution were recently studied by Alyami et al. (2023). For more recent studies, refer to Kayal and Balakrishnan (2023), Nair and Sathar (2024) and Smitha et al. (2024).

Weighted distributions provide a valuable tool for modeling statistical data in situations where standard distributions may not accurately capture the underlying characteristics of the data. Guiasu (1986) applied weighted entropy in order to balance the degree of homogeneity and information contained in a data partition into classes. The idea of weighted distributions was used in a number of domains, such as biostatistics (Wang, 1996), reliability modeling (Navarro et al., 2001) and renewal theory (Sunoj and Mayi, 2006). In the context of theoretical neurobiology, uncertainty measures based on the concept of weighted entropy were explored by Belis and Guiasu (1968). Di Crescenzo and Longobardi (2006) extended the concept of weighted entropy to residual and past lifetimes, introducing weighted residual and past entropies. This work builds upon previous research by Belzunce et al. (2004) and Nanda and Paul (2006), which characterized distribution functions using weighted dynamic measures. Misagh and Yari (2011) further investigated this concept by studying the weighted differential information measure for two-sided truncated random variables. Sunoj and Linu (2012) introduced a new measure of uncertainty based on the length-biased weighted function and called it dynamic weighted CRRE (DWCRRE). Based on the SF given in Equation (1), the DWCRRE is as follows:

$$R^*(c) = (1-c)^{-1} \log\left(\left(\overline{F}(t)\right)^{-c} \int_t^\infty y\left(\overline{F}(y)\right)^c dy\right); \quad c > 0, c \neq 1,$$
(2)

where y is the length-biased weighted function. The dynamic weighted CRTE (DWCRTE), introduced by Khammar and Jahanshahi (2018), is another significant weighted metric. It is defined as:

$$T^*(c) = (c-1)^{-1} \left( 1 - \left(\overline{F}(t)\right)^{-c} \int_t^\infty y\left(\overline{F}(y)\right)^c dy \right); \quad c > 0, c \neq 1.$$
(3)

Using Type II right-censored data, a weighted version of CRTE and DCRTE was created by Khammar and Jahanshahi (2018), and many of its reliability properties.

#### 1.3. Work Motivation

As far as we are aware, no research so far has taken into account the PT-IIC for dynamic weighted cumulative residual in entropy measurements. Therefore, our research question is: "How to find the estimate of dynamic weighted cumulative residual entropy measures under PT-IIC for the Burr XII distribution (BXIID)?". We decided to investigate this topic because of the significance of the BXIID and its widespread application in numerous sectors (as shown in Section 2). Our work involves the following steps:

- The DWCRE based on the Havrda and Charvat measure is defined following the idea of Sunoj and Linu (2012). This measure is called the dynamic weighted cumulative residual Havrda and Charvat entropy (DWCRHCE).
- The estimators for the DWCRRE, DWCRTE, and DWCRHCE are derived using both Bayesian and non-Bayesian techniques. Both symmetric and asymmetric loss

functions yield Bayesian estimators for the suggested measures. Also, because there are no explicit forms for the BEs of various measures, we use the Markov chain Monte Carlo (MCMC) approach to approximate the estimates.

• Simulation studies are employed to evaluate and contrast the precision of various approximations about their mean squared error (MSE) and average of estimates. For illustration reasons, application to real data is shown.

The rest of the paper is organized in the following way: Section 2 presents a model description. Dynamic weighted cumulative residual entropy expressions are determined in Section 3. The PT-IIC and maximum likelihood estimation are used in Section 4. The BE of the DWCRE for the BXIID under symmetric and asymmetric loss functions is presented in Section 5. Section 6 provides an example of a real-data application. A simulation study is provided in Section 7. Based on the outcomes of the numerical studies, the paper draws a few conclusions in the last section.

#### 2. Model Description

The Burr distribution is a versatile family that covers several widely used distributions as limiting asymptotic approximations, and it comprises a broad range of distribution shapes. A significant amount of the curve shape properties in the Pearson family are covered by correctly selecting the parameters of the Burr distribution, as Burr (1942) showed. Because its shape parameter generates a variety of forms that are excellent fits for different data, the BXIID has been used in research related to medical, business, chemical engineering, quality control and reliability. Evans and Simons (1975), Wingo (1993) and Gupta et al. (1996) are a few references to consult for a detailed explanation of such circumstances. The PDF and SF of the BXIID have the following specifications:

$$f(y) = \delta \lambda y^{\delta - 1} \left( 1 + y^{\delta} \right)^{-(\lambda + 1)}, y > 0,$$
(4)

$$\overline{F}(y) = \left(1 + y^{\delta}\right)^{-\lambda}, y > 0,$$
(5)

where  $\delta > \text{and} \lambda > 0$  are shape parameters. Recently, numerous scholars have conducted extensive research on the estimation utilizing the BXIID. Estimation in stepstress partially accelerated life tests for the BXIID using Type I censoring was covered by Abd-Elfattah et al. (2008). Works on BXIID inferences under the PCS were discussed by, for example, Mousa and Jaheen (2002), Wu and Yu (2005), Soliman (2005), Li et al. (2007) and Hassan et al. (2024a). According to Panahi and Sayyareh (2014), statistical inference and prediction about BXIID parameters based on a Type II censored sample were covered. On the basis of the competing risk model, Qin and Gui (2020) derived the MLE and BE of BXIID parameters.

## 3. Dynamic Weighted Cumulative Residual Entropy Expressions

Inspired by the DWCRRE entropy developed by Sunoj and Linu (2012) and the DWCRTE presented by Khammar and Jahanshahi (2018), we introduce two novel information measures: the DCR Havrda and Charvat entropy (DCRHCE) and the second measure is the DWCRHCE. This measure is based on the cumulative residual entropy originally proposed by Rao et al. (2004), and the dynamic cumulative residual entropy proposed by Asadi and Zohrevand (2007) and Sati and Gupta (2015).

**Definition:** The DCRHCE and DWCRHCE of a random variable *Y* of order *c* are defined by:

$$H(c) = (2^{1-c} - 1)^{-1} ((\overline{F}(t))^{-c} \int_t^{\infty} (\overline{F}(y))^c - 1); c > 0, c \neq 1,$$
  

$$H^*(c) = (2^{1-c} - 1)^{-1} ((\overline{F}(t))^{-c} \int_t^{\infty} y(\overline{F}(y))^c - 1); c > 0, c \neq 1,$$
(6)

where *y*, is the length-biased weighted function. Now, the expressions of DWCRRE, DWCRTE, and DWCRHCE for the BXIID are obtained. To do so, we first obtain the following integral by using the SF given in Equation (5)

$$I = \int_{t}^{\infty} y(\bar{F}(y))^{c} dy = \int_{t}^{\infty} y(1+y^{\delta})^{-\lambda c} dy.$$
<sup>(7)</sup>

Using the transformation  $z = y^{\delta}$ ,  $y = (z)^{\frac{1}{\delta}}$ ,  $dy = \delta^{-1} z^{\frac{1}{\delta}-1} dz$  in Equation (7) yields:

$$I = \int_{t^{\delta}}^{\infty} z^{\frac{1}{\delta}} (1+z)^{-\lambda c} \frac{1}{\delta} z^{\frac{1}{\delta}-1} dz = \frac{1}{\delta} \int_{t^{\delta}}^{\infty} (1+z)^{-\lambda c} z^{\frac{2}{\delta}-1} dz.$$

Use the transformation  $x = (1 + z)^{-1}$ , then  $z = \frac{(1-x)}{x}$ , and  $dz = \frac{-dx}{x^2}$ , thus the integral *I* is as follows:

$$I = \frac{1}{\delta} \int_{0}^{(1+t^{\delta})^{-1}} x^{\lambda c} \left(\frac{1-x}{x}\right)^{\frac{2}{\delta}-1} \frac{dx}{x^{2}} = \frac{1}{\delta} \int_{0}^{(1+t^{\delta})^{-1}} x^{\lambda c-\frac{2}{\delta}-1} (1-x)^{\frac{2}{\delta}-1} dx = \frac{1}{\delta} B\left(\frac{2}{\delta}, \lambda c-\frac{2}{\delta}, \left(1+t^{\delta}\right)^{-1}\right),$$
(8)

where  $B(a, b, x) = \int_0^x y^{a-1}(1-y)^{b-1} dy$  is the incomplete beta function. Now the formula of the DWCRRE is obtained by inserting Equation (8) in Equation (2), in the following way:

$$R^*(c) = \frac{1}{(1-c)} log\left(\frac{1}{\delta(1+t^{\delta})^{-\lambda^c}} B\left(\frac{2}{\delta}, \lambda c - \frac{2}{\delta}, \left(1+t^{\delta}\right)^{-1}\right)\right).$$
(9)

Also, expressions of DWCRTE and DWCRHCE measures are obtained by inserting Equation (8) in Equations (3) and (6), respectively, as below:

$$T^{*}(c) = \frac{1}{(c-1)} \left[ 1 - \frac{1}{\delta(1+t^{\delta})^{-\lambda c}} \left( B\left(\frac{2}{\delta}, \lambda c - \frac{2}{\delta}, \left(1 + t^{\delta}\right)^{-1}\right) \right) \right],$$
(10)

$$H^{*}(c) = \frac{1}{(2^{1-c}-1)} \left[ \frac{1}{\delta(1+t^{\delta})^{-\lambda c}} \left( B\left(\frac{2}{\delta}, \lambda c - \frac{2}{\delta}, \left(1+t^{\delta}\right)^{-1}\right) \right) - 1 \right].$$
(11)

Note that expressions (9), (10) and (11) are a function of parameters  $\delta$ ,  $\lambda$ , and c.

### 4. Maximum Likelihood Estimators

Here, the MLEs of DWCRRE, DWCRTE, and DWCRHCE of the BXIID, based on PT-IIC samples, are obtained. Assume that  $y_{(1)} \le y_{(2)} \le ... \le y_{(m)}$  be the PT-IIC of

size *m* from a sample of size *n* taken from density given in Equation (4) and SF given in Equation (5) with censoring scheme  $r_1, r_2, ..., r_m$ . Based on the PT-IIC sample, the likelihood function reads:

$$L(\delta,\lambda) = D \prod_{i=1}^{m} f(y_{(i)}) \left[ 1 - F(y_{(i)}) \right]^{r_i} = D \delta^m \lambda^m \prod_{i=1}^{m} y_{(i)}^{\delta-1} \left( 1 + y_{(i)}^{\delta} \right)^{-\lambda(r_i+1)-1},$$

where  $D = n(n - r_1 - 1)(n - r_1 - r_2 - 2)...n - m + 1 - \sum_{i=1}^{m-1} r_i$ . Additionally, the corresponding log-likelihood function, say  $\ell^*$ , is

 $\ell^* \propto m \ln(\delta) + m \ln(\lambda) + (\delta - 1) \sum_{i=1}^m \ln(y_{(i)}) - \sum_{i=1}^m [\lambda(r_i + 1) + 1] \ln(1 + y_{(i)}^{\delta}).$ 

Then, the first derivative of the log- likelihood function, with respect to the parameters  $\delta$  and  $\lambda$ , are as follows:

$$\frac{\partial \ell^*}{\partial \delta} = \frac{m}{\delta} + \sum_{i=1}^m ln y_{(i)} - \sum_{i=1}^m \frac{[\lambda(r_i+1)+1] \ln y_{(i)}}{(1+y_{(i)}^{-\delta})},$$
(12)

and

$$\frac{\partial \ell^*}{\partial \lambda} = \frac{m}{\lambda} - \sum_{i=1}^m (r_i + 1) \ln(1 + y_{(i)}^{\delta}).$$
(13)

The MLEs of  $\delta$  and  $\lambda$  are determined by solving Equations (12) and (13) after simultaneously setting them to zero by using a numerical technique such as the Newton-Raphson method, to get  $\hat{\delta}_{ML}$  and  $\hat{\lambda}_{ML}$ . Based on the invariance property, the MLEs of DWCRRE, DWCRTE, and DWCRHCE are obtained by inserting  $\hat{\delta}_{ML}$  and  $\hat{\lambda}_{ML}$ in Equations (9), (10), and (11).

#### 5. Bayesian Estimator

This part covers the BE of  $T^*(c)$ ,  $R^*(c)$  and  $H^*(c)$  under both symmetric and asymmetric loss functions for the two-parameter BXIID. It is assumed that the prior of parameters  $\delta$  and  $\lambda$  have independent gamma priors. The joint prior distribution can be written as:

$$\pi_0(\delta,\lambda) \propto \delta^{b_1-1} \lambda^{b_2-1} e^{-(\delta a_1+\lambda a_2)}$$

where the hyper-parameters  $a_1, a_2, b_1$ , and  $b_2$  are known and non-negative. The gamma prior was chosen for its flexibility in modeling a wide range of prior beliefs. They may take on a wide range of forms based on hyper-parameters. The joint posterior for parameters, denoted by  $\pi(\delta, \lambda)$ , is

$$\pi(\delta,\lambda) = W^{-1}\delta^{m+b_1-1}\lambda^{m+b_2-1}e^{-(\delta a_1+\lambda a_2)}\prod_{i=1}^m y_{(i)}^{\delta-1}(1+y_{(i)}^{\delta})^{-[\lambda(r_i+1)+1]},$$
$$W = \int_0^\infty \int_0^\infty \delta^{m+b_1-1}\lambda^{m+b_2-1}e^{-(\delta a_1+\lambda a_2)}\prod_{i=1}^m y_{(i)}^{\delta-1}(1+y_{(i)}^{\delta})^{-[\lambda(r_i+1)+1]}d\delta d\lambda.$$

where

So, the conditional posterior distribution of the unknown parameters  $\delta$  and  $\lambda$ , is given, respectively, by:

$$\pi^*\left(\delta\left|\lambda,\underline{y}\right) \propto \delta^{m+b_1-1} e^{-\delta\left(a_1+\sum_{l=1}^m lny_{(l)}\right) - \sum_{l=1}^m [\lambda(r_l+1)+1] ln(1+y_{(l)}^{\delta})},\tag{14}$$

and

$$\boldsymbol{\pi}^{**}\left(\boldsymbol{\lambda} \left| \boldsymbol{\delta}, \underline{\boldsymbol{y}} \right) \propto \text{Gamma}\left( m + b_2, a_2 + \sum_{i=1}^{m} (r_i + 1) \ln(1 + y_{(i)}^{\delta}) \right).$$
(15)

The Bayesian estimators of  $g^{\bullet}(c)$ , under squared error loss function (SLF), linear exponential loss function (LLF), and precautionary loss function (PRLF) are obtained in the following way:

$$\widehat{g}_{SLF}^{\bullet}(c) = E\left[g^{\bullet}(c) \left| \frac{y}{y} \right] = W^{-1} \int_{0}^{\infty} \int_{0}^{\infty} g^{\bullet}(c) \delta^{m+b_{1}-1} \lambda^{m+b_{2}-1} e^{-(\delta a_{1}+\lambda a_{2})} \prod_{i=1}^{m} y_{(i)}^{\delta-1} \left(1 + y_{(i)}^{\delta}\right)^{-[\lambda(r_{i}+1)+1]} d\delta d\lambda,$$
(16)

$$\widehat{g}_{LLF}^{\bullet}(c) = \frac{-1}{\tau} ln \left[ E \left( e^{-\tau g^{\bullet}(c)} \left| \underline{y} \right) \right] = W^{-1} \int_{0}^{\infty} \int_{0}^{\infty} \delta^{m+b_{1}-1} \lambda^{m+b_{2}-1} e^{-(\tau g^{\bullet}(c)+\delta a_{1}+\lambda a_{2})} \prod_{i=1}^{m} y_{(i)}^{\delta-1} \left( 1 + y_{(i)}^{\delta} \right)^{-[\lambda(r_{i}+1)+1]} d\delta d\lambda, \quad (17)$$

and

$$\widehat{g}_{PRLF}^{\bullet}(c) = \left[ E\left( (g^{\bullet}(c))^2 \left| \underline{y} \right) \right]^{0.5} = W^{-1} \int_0^\infty \int_0^\infty (g^{\bullet}(c))^2 \delta^{m+b_1-1} \lambda^{m+b_2-1} e^{-(\delta a_1+\lambda a_2)} \prod_{i=1}^m y_{(i)}^{\delta-1} (1+y_{(i)}^{\delta})^{-[\lambda(r_i+1)+1]} d\delta d\lambda,$$
(18)

where  $g^{\bullet}(c) = R^{*}(c)$  to obtain the DWCRRE,  $g^{\bullet}(c) = T^{*}(c)$  to calculate the DWCRTE, and  $g^{\bullet}(c) = H^{*}(c)$  to produce the DWCRHCE. The Gibbes sampler, Metropolis-Hastings (M-H), and random walk Metropolis algorithms are used to generate the MCMC samples from the posterior density functions (14) and (15), respectively, because integrals (16)–(18) do not take a closed form. As a result, the BEs of  $\delta$  and  $\lambda$  under the SLF, LLF, and PRLF are calculated from their posteriors as the mean of the simulated samples. The M-H algorithm is one of the most famous subclasses of the MCMC method in Bayesian literature. It is used to simulate the deviates from the posterior density and produce good approximate results. The M-H algorithm uses an acceptance/rejection rule to converge to the target distribution. The initial values of the unknown parameters  $(\delta, \lambda)$  must be specified, along with a suggested distribution, in order to implement the M-H algorithm for the DWCRTE, DWCRRE, and the DWCRHCE of the BXIID. A normal distribution will be used to calculate the proposal distribution, i.e.,  $h(\delta'|\delta)$  and  $h(\lambda'|\lambda) \equiv N(\theta, \sigma_{\theta}^2)$ , where  $\theta \equiv (\delta, \lambda)$  and  $\sigma_{\theta}^2$  is the variance-covariance matrix (Va-CoM) for the MLEs of  $(\delta, \lambda)$ . The MLE for  $\theta$  is taken into account, that is, $\theta^{(0)} = \hat{\theta}_{MLE}$ . Asymptotic Va-CoM, say  $I^{-1}(\hat{\theta}_{MLE})$ , where I(.) is the Fisher information matrix, and is assumed to represent the choice of  $\sigma_{\theta}^2$ . First, the M-H algorithm employs the steps mentioned below to extract a sample from the posterior density given by Equations (14) and (15).

**Step 1:** Set the initial value of  $\theta$  as  $\theta^{(0)} = \hat{\theta}_{MLE}$ .

**Step 2:** For i = 1, 2, 3, ..., M repeat the following steps:

2.1: Set  $\theta = \theta^{(i-1)}$ .

2.2: Generate $\lambda_1^i$  from Gamma $(m + b_2, a_2 + \sum_{i=1}^m \lambda(r_i + 1) \ln(1 + y_{(i)}^{\delta}))$ .

- 2.3: Create a new candidate parameter value using  $N(\theta, S_{\theta})$ .
- 2.4: Compute the formula  $\phi = \frac{\pi(\theta'|y)}{\pi(\theta|y)}$  where  $\pi(.)$  is the posterior density of Equations (14) and (15).
- 2.5: Create a sample u from the uniform U(0,1) distribution.

2.6: Accept or reject the new candidate 
$$\theta'$$
. 
$$\begin{cases} Ifu \le \phi put \theta^{(i)} = \theta' \\ elsewhereput \theta^{(i)} = \theta. \end{cases}$$

**Step 3:** Obtain the Bayesian estimator of  $\theta$  and compute the DWCRTE, DWCRRE, and DWCRHCE functions of  $T^*(c)$ ,  $R^*(c)$  and  $H^*(c)$  with respect to the loss functions as follows:

$$\begin{split} \hat{H}^{*}(c) &= \frac{1}{M-Q} \sum_{i=Q+1}^{M} H^{*}(c,\theta^{(i)}), \hat{R}^{*}(c) = \frac{1}{M-Q} \sum_{i=Q+1}^{M} R^{*}(c,\theta^{(i)}), \hat{T}^{*}(c) \\ &= \frac{1}{M-Q} \sum_{i=Q+1}^{M} T^{*}(c,\theta^{(i)}), \end{split}$$

where *Q* is the number of samples that have been burned. Ultimately, the estimates of DWCRTE, DWCRRE, and DWCRHCE are obtained by subtracting 2,000 burn-in samples from the 10,000 samples that the posterior density produced.

#### 6. Real Data Analysis

The data set represents the monthly water capacity data from the Shasta reservoir in California, USA, and was taken for the month of February from 1991 to 2010 (http://cdec.water.ca.gov/reservoir\_map.html). The maximum capacity of the reservoir is 4552000 AF, and the data set was transformed to the interval [0, 1] (for more details, see Nadar et al., 2013). In the first step, it should be checked if the BXIID is well fitted to this data. By fitting BXIID, the Kolmogorov-Smirnov distance and associated p-value are 7.7852, 7.8451, 0.22479 and 0.2274, respectively. The p-values show that BXIID yields suitable fits for the given dataset. In Figure 1, the empirical distribution functions and histogram are provided.





The number of stages in the PT-IIC scheme is assumed to be m = 12 and the removed items  $r_i$  are assumed as in the following scheme (Sch.):

 $S_1: (8,0^{*11}), S_2: (4,0^{*10}, 4), S_3: (0^{*2}, 1^{*8}, 0^{*2}), and S_4: (0^{*11}, 8).$ 

The complete case also considers instances where n = m = 2 and  $r_i = 0$ , i = 1, 2, ..., n. Table 1 uses the PT-IIC to compute MLEs based on the produced data for each censoring strategy. Then employed to compute  $T^*(c)$ ,  $R^*(c)$  and  $H^*(c)$  given t and c, where t = 0.1, 0.2, and c = 1.5, 2.5. For the BEs, the M-H algorithm will be used under different loss functions in the case of a uniform prior, where  $a_1 = a_2 = b_1 = b_2 = 0.001$ . Different loss functions, including SLF, LLF-1( $\tau = 0.5$ ), LLF-2( $\tau = -0.5$ ) and PRLF are assumed. After that, the estimated values are calculated using the previous values. The estimates of DWCRTE, DWCRRE, and DWCRHCE are then obtained after 2,000 burn-in samples are subtracted from the 10,000 samples that the posterior density produced.

Convergence of the MCMC estimates for the DWCRTE, DWCRRE, and the DWCRHCE using M-H algorithms is shown in two figures in the case of complete sampling. Each figure shows the plot, histogram and cumulative mean where t = 0.1



and c = 1.5 in Figure 2 while Figure 3 represents the case where t = 0.2 and c = 2.5.

**Figure 2**. MCMC convergence of DWCRTE, DWCRRE, and DWCRHCE estimates at t = 0.1 and c = 1.5*Source: created by researchers utilizing the R programming language.* 



**Figure 3.** MCMC convergence of DWCRTE, DWCRRE, and DWCRHCE estimates at t = 0.2 and c=2.5Source: created by researchers utilizing the *R* programming language.

Sch	Mathad	1	t = 0.1, c =	1.5		t = 0.2, c = 2	.5
Scn.	Method	DWCRTE	DWCRRE	DWCRHCE	DWCRTE	DWCRRE	DWCRHCE
	MLE	1.5100	2.8129	2.5777	0.5322	1.0673	1.2349
	SLF	1.5249	2.8747	2.6032	0.5406	1.1102	1.2544
<i>S</i> <sub>1</sub>	LLF-1	1.5147	2.8321	2.5857	0.5401	1.1074	1.2531
	LLF-2	1.5612	3.0335	2.6651	0.5488	1.1550	1.2733
	PRLF	1.5309	2.9001	2.6134	0.5417	1.1159	1.2568
	MLE	1.4986	2.7671	2.5583	0.5427	1.1217	1.2594
<i>S</i> <sub>2</sub>	SLF	1.4860	2.7175	2.5368	0.5414	1.1147	1.2563
	LLF-1	1.4828	2.7049	2.5313	0.5417	1.1162	1.2569
	LLF-2	1.4915	2.7388	2.5461	0.5415	1.1151	1.2565
	PRLF	1.4939	2.7484	2.5503	0.5431	1.1235	1.2601
	MLE	1.4282	2.5043	2.4381	0.5105	0.9674	1.1845
	SLF	1.4396	2.5446	2.4576	0.5171	0.9964	1.1999
<b>S</b> <sub>3</sub>	LLF-1	1.4350	2.5281	2.4497	0.5185	1.0025	1.2031
	LLF-2	1.5249	2.8747	2.6031	0.5375	1.0939	1.2471
	PRLF	1.4561	2.6043	2.4858	0.5209	1.0136	1.2087
	MLE	1.3820	2.3487	2.3591	0.4951	0.9047	1.1487
	SLF	1.3866	2.3638	2.3671	0.4998	0.9234	1.1597
<b>S</b> <sub>4</sub>	LLF-1	1.3971	2.3981	2.3849	0.5059	0.9484	1.1740
	LLF-2	1.3761	2.3298	2.3491	0.4932	0.8976	1.1445
	PRLF	1.3922	2.3822	2.3767	0.5006	0.9267	1.1616
	MLE	1.5228	2.8660	2.5996	0.5384	1.0986	1.2492
	SLF	1.5324	2.9066	2.6160	0.5434	1.1256	1.2610
nlete	LLF-1	1.5272	2.8844	2.6071	0.5436	1.1265	1.2614
piece	LLF-2	1.5560	3.0104	2.6563	0.5485	1.1536	1.2728
	PRLF	1.5354	2.9195	2.6211	0.5438	1.1277	1.2619

Table 1. Estimates of different entropy measures under PT-IIC schemes given real dataset wherem=15

# 7. Simulation Study

This section evaluates the performance of estimate methodologies for BXIID under the PT-IIC scheme using a Monte Carlo simulation analysis. Specifically, maximum likelihood and Bayesian processes employing MCMC are considered. From the BXIID, we generate 1,000 random samples using the following guidelines:

- 1. Two cases of parameters of BXIID are assumed, namely:  $(\delta, \lambda) = (1.5, 2.5)$ and  $(\delta, \lambda) = (2.5, 1.5)$
- 2. Two cases of parameters of weighted entropy measures are assumed, namely: (t, c) = (0.5, 1.5) and (t, c) = (1.5, 2.5).
- 3. The true values of different entropy measures are listed in Table 2.

δ	λ	t	С	$T^{*}(c)$	$R^*(c)$	$H^*(c)$
1.5	2.5	0.5	1.5	1.3621	2.2853	2.3252
		1.5	2.5	0.3434	0.4825	0.7968
2.5	1.5	0.5	1.5	1.2567	1.9796	2.1453
		1.5	2.5	0.3943	0.5964	0.9115

Table 2. True values of different entropy measures

- 4. The sample size is assumed to be n = 40 and n = 60.
- 5. The number of stages in the PT-IIC scheme is m = 20, 30 at n = 40, m = 40 and 50 at n = 60.
- 6. Removed items  $r_i$  are assumed to have *n* and *m* values as shown in Table 3, where  $r_m = n m \sum_{i=1}^{m-1} r_i$  and  $r_i$  is the number of failure items.

Based on the produced data and the previously-made assumptions, the MLEs are calculated using PT-IIC. After that, the MLEs are used to calculate  $T^*(c)$ ,  $R^*(c)$  and  $H^*(c)$  given the values of *t* and *c*. For the Bayesian method, BEs using the M-H algorithm under different loss functions in the case of gamma prior are computed, where the following hyper-parameters are assumed:  $(a_1, b_1, a_2, b_2) = (1.5, 2.5, 1.75, 2.75)$ .

Different loss functions, including SLF, LLF-1( $\tau = 0.5$ ), LLF-2( $\tau = -0.5$ ) and PRLF, are given. These values are then employed to determine the estimated values. The average (Avg.) of all entropy estimates and the Avg. of the MSE are presented in Tables 4 (a) to 5 (d), which incorporate all of the Monte Carlo simulation's inputs.

( <i>n</i> , <i>m</i> )	Censoring Schemes								
	$S_1$	$S_2$	S <sub>3</sub>	$S_4$					
(40,20)	$(20, 0^{*19})$	$(10, 0^{*18}, 10)$	$(1^{*20})$	(0*19, 20)					
(40,30)	$(10, 0^{*29})$	$(5, 0^{*28}, 5)$	$(0^{*10}, 1^{*10}, 0^{*10})$	$(0^{*29}, 10)$					
(60,40)	$(20, 0^{*39})$	$(10, 0^{*38}, 10)$	$(0^{*10}, 1^{*20}, 0^{*10})$	$(0^{*39}, 20)$					
(60,50)	$(10, 0^{*49})$	$(5, 0^{*48}, 5)$	$(0^{*20}, 1^{*10}, 0^{*20})$	$(0^{*49}, 10)$					

Table 3. Numerous patterns for removing items from life test at different number of stages

Here,  $(2^{*4}, 0)$ , for example, means that the censoring scheme employed is (2,2,2,2,0). *Source: created by researchers.* 

**Table 4 (a).** The Avg. and MSE of different weighted entropy estimates for BXIID under PT-IIC schemes at( $\delta$ ,  $\lambda$ ) = (1.5, 2.5), and (n, m) = (40, 20)

( <i>t</i> , <i>c</i> )	Sch.	Estimate	DWCRTE		DWCRRE		DWCRHCE	
			Avg.	MSE	Avg.	MSE	Avg.	MSE
	<i>S</i> <sub>1</sub>	MLE	1.4108	0.0794	2.7101	2.6286	2.4084	0.2314
(0 5 1 5)		SLF	0.9446	0.4437	1.4681	1.3845	1.6126	1.2929
(0.5,1.5)		LLF-1	0.8564	0.5946	1.3158	1.6815	1.4620	1.7328
		LLF-2	1.0257	0.3293	1.6214	1.1383	1.7510	0.9596

(t, c) Sch.		Estimate	DWG	CRTE	DW	CRRE	DWCF	RHCE
(t, c)	Sch.	Estimate	Avg.	MSE	Avg.	MSE	Avg.	MSE
		PRLF	1.0293	0.3130	1.6195	1.1060	1.7571	0.9121
		MLE	1.3937	0.2089	3.0689	12.2305	2.3792	0.6087
		SLF	0.4167	3.2758	0.8684	3.3052	0.7114	9.5464
	<i>S</i> <sub>2</sub>	LLF-1	0.2216	4.7021	0.6644	3.9941	0.3783	13.7029
		LLF-2	0.5888	2.2816	1.0751	2.7086	1.0051	6.6492
		PRLF	0.6137	2.0044	1.0844	2.6074	1.0477	5.8414
		MLE	1.4055	0.1730	3.0630	10.8346	2.3994	0.5042
		SLF	0.4537	2.7111	0.8827	3.1863	0.7746	7.9007
	$S_3$	LLF-1	0.2675	3.8704	0.6797	3.8568	0.4566	11.2793
		LLF-2	0.6184	1.9003	1.0883	2.6073	1.0557	5.5378
		PRLF	0.6402	1.6882	1.0959	2.5141	1.0929	4.9198
		MLE	1.4494	0.1353	3.6653	12.6546	2.4743	0.3944
		SLF	0.7864	0.4795	1.1076	1.8434	1.3424	1.3972
	$S_4$	LLF-1	0.7149	0.5654	0.9811	2.1068	1.2205	1.6476
		LLF-2	0.8485	0.4207	1.2309	1.6438	1.4485	1.2260
		PRLF	0.8377	0.4300	1.2074	1.6717	1.4300	1.2532
		MLE	0.3752	0.0147	0.6654	0.6670	0.8707	0.0792
		SLF	0.2018	0.0438	0.2752	0.0897	0.4682	0.2358
	<i>S</i> <sub>1</sub>	LLF-1	0.1756	0.0545	0.2384	0.1057	0.4076	0.2936
		LLF-2	0.2271	0.0349	0.3130	0.0763	0.5269	0.1880
		PRLF	0.2281	0.0337	0.3130	0.0742	0.5292	0.1816
		MLE	0.3793	0.0274	0.8829	2.5356	0.8800	0.1473
		SLF	0.0796	0.1314	0.1351	0.1853	0.1846	0.7077
	$S_2$	LLF-1	0.0394	0.1632	0.0905	0.2174	0.0913	0.8786
		LLF-2	0.1184	0.1049	0.1815	0.1565	0.2747	0.5647
(1525)		PRLF	0.1223	0.0994	0.1836	0.1518	0.2839	0.5350
(1.5,2.5)		MLE	0.3818	0.0250	0.8531	2.2068	0.8859	0.1347
		SLF	0.0848	0.1228	0.1380	0.1797	0.1968	0.6611
	<b>S</b> <sub>3</sub>	LLF-1	0.0453	0.1526	0.0937	0.2113	0.1051	0.8215
		LLF-2	0.1230	0.0979	0.1843	0.1516	0.2855	0.5271
		PRLF	0.1265	0.0930	0.1860	0.1471	0.2936	0.5008
		MLE	0.4105	0.0254	1.1969	4.9892	0.9526	0.1366
		SLF	0.1754	0.0390	0.2198	0.0918	0.4071	0.2098
	$S_4$	LLF-1	0.1592	0.0436	0.1955	0.1015	0.3694	0.2348
		LLF-2	0.1901	0.0358	0.2436	0.0849	0.4412	0.1926
		PRLF	0.1878	0.0360	0.2394	0.0855	0.4358	0.1941

(4.)	6.1		DWG	CRTE	DWG	CRRE	DWCRHCE	
(t, c)	Sch.	Estimate	Avg.	MSE	Avg.	MSE	Avg.	MSE
		MLE	1.4217	0.0503	2.6143	0.6428	2.4270	0.1465
		SLF	1.1547	0.1414	1.8470	0.6809	1.9712	0.4122
	<i>S</i> <sub>1</sub>	LLF-1	1.1107	0.1751	1.7480	0.7857	1.8961	0.5102
		LLF-2	1.1966	0.1143	1.9466	0.5968	2.0427	0.3330
		PRLF	1.1964	0.1113	1.9420	0.5834	2.0424	0.3243
		MLE	1.4242	0.0626	2.6550	0.7982	2.4313	0.1825
		SLF	1.0862	0.2244	1.7193	0.9094	1.8543	0.6540
	<b>S</b> <sub>2</sub>	LLF-1	1.0335	0.2789	1.6101	1.0545	1.7643	0.8128
		LLF-2	1.1362	0.1802	1.8293	0.7898	1.9397	0.5252
		PRLF	1.1370	0.1744	1.8256	0.7714	1.9411	0.5082
(0.5,1.5)		MLE	1.4242	0.0601	2.6487	0.7695	2.4313	0.1752
		SLF	1.0926	0.2134	1.7294	0.8854	1.8653	0.6219
	<b>S</b> <sub>3</sub>	LLF-1	1.0415	0.2643	1.6224	1.0243	1.7779	0.7703
		LLF-2	1.1412	0.1719	1.8373	0.7709	1.9482	0.5011
		PRLF	1.1416	0.1668	1.8327	0.7538	1.9488	0.4861
		MLE	1.4242	0.0785	2.7197	1.5945	2.4313	0.2287
		SLF	1.0046	0.3581	1.5816	1.1984	1.7149	1.0435
	<i>S</i> <sub>4</sub>	LLF-1	0.9404	0.4489	1.4612	1.3958	1.6054	1.3081
		LLF-2	1.0651	0.2848	1.7036	1.0321	1.8182	0.8300
		PRLF	1.0673	0.2734	1.7008	1.0069	1.8221	0.7967
		MLE	0.3770	0.0103	0.6030	0.1413	0.8747	0.0554
	<i>S</i> <sub>1</sub>	SLF	0.2679	0.0189	0.3700	0.0496	0.6217	0.1016
		LLF-1	0.2525	0.0224	0.3444	0.0556	0.5859	0.1204
		LLF-2	0.2830	0.0160	0.3960	0.0449	0.6567	0.0859
		PRLF	0.2831	0.0156	0.3954	0.0439	0.6570	0.0839
		MLE	0.3799	0.0127	0.6378	0.3206	0.8815	0.0685
		SLF	0.2445	0.0271	0.3361	0.0637	0.5674	0.1457
	<i>S</i> <sub>2</sub>	LLF-1	0.2270	0.0321	0.3085	0.0719	0.5266	0.1728
		LLF-2	0.2617	0.0228	0.3643	0.0571	0.6072	0.1227
(1525)		PRLF	0.2621	0.0222	0.3639	0.0558	0.6081	0.1195
(1.3,2.3)		MLE	0.3792	0.0121	0.6334	0.3127	0.8799	0.0651
		SLF	0.2468	0.0259	0.3388	0.0619	0.5726	0.1392
	<b>S</b> <sub>3</sub>	LLF-1	0.2297	0.0306	0.3118	0.0697	0.5330	0.1649
		LLF-2	0.2635	0.0218	0.3664	0.0555	0.6114	0.1175
		PRLF	0.2637	0.0213	0.3657	0.0543	0.6118	0.1147
		MLE	0.3821	0.0155	0.7030	0.7839	0.8865	0.0836
		SLF	0.2189	0.0380	0.3008	0.0804	0.5079	0.2043
	<i>S</i> <sub>4</sub>	LLF-1	0.1988	0.0451	0.2711	0.0913	0.4614	0.2429
		LLF-2	0.2385	0.0318	0.3314	0.0713	0.5534	0.1714
		PRLF	0.2392	0.0309	0.3312	0.0696	0.5550	0.1664

**Table 4 (b).** Avg. and MSE of different weighted entropy estimates for BXIID under PT-IIC schemes at  $(\delta, \lambda) = (1.5, 2.5)$  and (n, m) = (40, 30)

(4 )	Sch.	E-thurst.	DWC	CRTE	DWG	CRRE	DWCRHCE	
(1, 0)	Sch.	Estimate	Avg.	MSE	Avg.	MSE	Avg.	MSE
		MLE	1.4210	0.0362	2.5731	0.4596	2.4258	0.1055
	<i>S</i> <sub>1</sub>	SLF	1.2338	0.0707	2.0056	0.4220	2.1062	0.2060
		LLF-1	1.2049	0.0839	1.9326	0.4710	2.0569	0.2445
		LLF-2	1.2617	0.0598	2.0792	0.3841	2.1539	0.1744
		PRLF	1.2606	0.0589	2.0740	0.3773	2.1520	0.1715
		MLE	1.4226	0.0486	2.6113	0.6132	2.4285	0.1415
		SLF	1.1639	0.1287	1.8595	0.6310	1.9870	0.3751
	<i>S</i> <sub>2</sub>	LLF-1	1.1274	0.1540	1.7750	0.7132	1.9246	0.4487
		LLF-2	1.1992	0.1076	1.9450	0.5639	2.0472	0.3137
$(0 \in 1 \in )$		PRLF	1.1986	0.1054	1.9400	0.5537	2.0461	0.3072
(0.3,1.3)		MLE	1.4234	0.0457	2.6074	0.5854	2.4300	0.1331
		SLF	1.1741	0.1171	1.8784	0.5962	2.0043	0.3413
	<b>S</b> <sub>3</sub>	LLF-1	1.1392	0.1396	1.7965	0.6718	1.9448	0.4069
		LLF-2	1.2078	0.0983	1.9612	0.5346	2.0619	0.2865
		PRLF	1.2068	0.0966	1.9554	0.5257	2.0601	0.2814
		MLE	1.4209	0.0642	2.6426	0.7779	2.4256	0.1871
		SLF	1.0802	0.2275	1.7009	0.9062	1.8440	0.6629
	<i>S</i> <sub>4</sub>	LLF-1	1.0334	0.2748	1.6034	1.0341	1.7642	0.8007
		LLF-2	1.1251	0.1879	1.8001	0.7982	1.9207	0.5476
		PRLF	1.1253	0.1829	1.7953	0.7834	1.9209	0.5329
	<i>S</i> <sub>1</sub>	MLE	0.3734	0.0073	0.5727	0.0428	0.8663	0.0395
		SLF	0.2944	0.0107	0.4083	0.0323	0.6830	0.0576
		LLF-1	0.2836	0.0123	0.3890	0.0353	0.6579	0.0662
		LLF-2	0.3050	0.0094	0.4279	0.0301	0.7078	0.0505
		PRLF	0.3048	0.0092	0.4270	0.0296	0.7072	0.0497
		MLE	0.3757	0.0098	0.5874	0.0588	0.8717	0.0525
		SLF	0.2686	0.0176	0.3687	0.0463	0.6234	0.0948
	<i>S</i> <sub>2</sub>	LLF-1	0.2557	0.0204	0.3469	0.0512	0.5932	0.1096
		LLF-2	0.2815	0.0153	0.3909	0.0423	0.6531	0.0822
(1525)		PRLF	0.2814	0.0150	0.3901	0.0415	0.6529	0.0807
(1.3,2.3)		MLE	0.3754	0.0091	0.5843	0.0550	0.8710	0.0489
		SLF	0.2725	0.0161	0.3740	0.0436	0.6323	0.0869
	$S_3$	LLF-1	0.2601	0.0186	0.3529	0.0481	0.6035	0.1002
		LLF-2	0.2848	0.0140	0.3954	0.0400	0.6608	0.0755
		PRLF	0.2845	0.0138	0.3944	0.0393	0.6602	0.0743
		MLE	0.3771	0.0125	0.6227	0.2506	0.8749	0.0673
		SLF	0.2405	0.0272	0.3275	0.0631	0.5580	0.1466
	<i>S</i> <sub>4</sub>	LLF-1	0.2249	0.0316	0.3031	0.0705	0.5219	0.1704
		LLF-2	0.2558	0.0234	0.3527	0.0568	0.5937	0.1260
		PRLF	0.2559	0.0229	0.3519	0.0558	0.5939	0.1234

**Table 4(c)**. The Avg. and MSE of different weighted entropy estimates for BXIID under PT-IIC schemes at  $(n, (\delta, \lambda) = (2.5, 1.5) \text{ and } (n, m) = (60, 40)$ 

(4 )	C -L	F-45	DWCRTE		DWC	CRRE	DWCRHCE	
(t, c)	Scn.	Estimate	Avg.	MSE	Avg.	MSE	Avg.	MSE
		MLE	1.4218	0.0295	2.5565	0.3722	2.4272	0.0861
		SLF	1.2762	0.0462	2.1027	0.3103	2.1787	0.1347
	<i>S</i> <sub>1</sub>	LLF-1	1.2546	0.0531	2.0441	0.3366	2.1417	0.1549
		LLF-2	1.2973	0.0405	2.1615	0.2912	2.2147	0.1180
		PRLF	1.2962	0.0400	2.1569	0.2864	2.2128	0.1165
		MLE	1.4231	0.0332	2.5708	0.4189	2.4295	0.0966
		SLF	1.2558	0.0580	2.0556	0.3618	2.1439	0.1690
	<b>S</b> <sub>2</sub>	LLF-1	1.2323	0.0670	1.9938	0.3952	2.1037	0.1953
		LLF-2	1.2787	0.0505	2.1177	0.3365	2.1829	0.1472
(0 = 1 = 1)		PRLF	1.2777	0.0498	2.1131	0.3310	2.1812	0.1451
(0.5,1.5)		MLE	1.4231	0.0326	2.5692	0.4127	2.4293	0.0951
		SLF	1.2570	0.0570	2.0581	0.3586	2.1459	0.1661
	<b>S</b> <sub>3</sub>	LLF-1	1.2339	0.0657	1.9972	0.3912	2.1064	0.1916
		LLF-2	1.2796	0.0497	2.1194	0.3338	2.1844	0.1449
		PRLF	1.2784	0.0491	2.1145	0.3285	2.1824	0.1429
		MLE	1.4237	0.0373	2.5833	0.4680	2.4303	0.1087
		SLF	1.2340	0.0726	2.0070	0.4215	2.1066	0.2116
	<b>S</b> <sub>4</sub>	LLF-1	1.2083	0.0843	1.9414	0.4635	2.0627	0.2457
		LLF-2	1.2591	0.0628	2.0732	0.3886	2.1494	0.1830
		PRLF	1.2581	0.0618	2.0686	0.3822	2.1477	0.1802
		MLE	0.3726	0.0060	0.5657	0.0341	0.8646	0.0323
	<i>S</i> <sub>1</sub>	SLF	0.3098	0.0074	0.4331	0.0246	0.7189	0.0401
		LLF-1	0.3014	0.0083	0.4174	0.0262	0.6993	0.0449
		LLF-2	0.3181	0.0067	0.4490	0.0235	0.7382	0.0362
		PRLF	0.3179	0.0066	0.4482	0.0231	0.7377	0.0357
		MLE	0.3737	0.0068	0.5710	0.0391	0.8671	0.0364
		SLF	0.3017	0.0091	0.4198	0.0284	0.7001	0.0489
	<i>S</i> <sub>2</sub>	LLF-1	0.2927	0.0102	0.4034	0.0304	0.6792	0.0549
		LLF-2	0.3106	0.0081	0.4365	0.0269	0.7208	0.0438
(1525)		PRLF	0.3104	0.0080	0.4358	0.0265	0.7203	0.0432
(1.3,2.3)		MLE	0.3734	0.0066	0.5698	0.0380	0.8665	0.0355
		SLF	0.3023	0.0089	0.4207	0.0279	0.7015	0.0478
	<b>S</b> <sub>3</sub>	LLF-1	0.2935	0.0100	0.4045	0.0299	0.6810	0.0536
		LLF-2	0.3110	0.0080	0.4370	0.0265	0.7217	0.0429
		PRLF	0.3108	0.0079	0.4362	0.0260	0.7211	0.0423
		MLE	0.3744	0.0076	0.5757	0.0441	0.8688	0.0407
		SLF	0.2934	0.0110	0.4065	0.0325	0.6807	0.0591
	<i>S</i> <sub>4</sub>	LLF-1	0.2837	0.0124	0.3892	0.0351	0.6582	0.0667
		LLF-2	0.3029	0.0098	0.4241	0.0306	0.7029	0.0526
		PRLF	0.3028	0.0096	0.4234	0.0301	0.7025	0.0518

**Table 4(d).** The Avg. and MSE of different weighted entropy estimates for BXIID under PT-IIC schemes at  $(\delta, \lambda) = (2.5, 1.5)$  and (n, m) = (60, 50)

Estimate

MLE

SLF

Sch.

(*t*, *c*)

ghted er	ntropy est	imates for	BXIID un	der PT-IIC								
m) = (40, 20)												
TE DWCRRE DWCRHCE												
MSE	Avg.	MSE	Avg.	MSE								
0.0516	2.2314	0.7609	2.2307	0.1504								
0.1430	1.5868	0.5515	1.7901	0.4168								
0.1856	1.4846	0.6601	1.7007	0.5410								
0.1101	1.6882	0.4660	1.8738	0.3209								
0.1024	1.7018	0.4431	1.8873	0.2983								
0.0780	2.3729	3.1072	2.2422	0.2274								
0.4090	1.2871	1.0673	1.4748	1.1918								
0.5359	1.1649	1.2812	1.3432	1.5618								
0.3116	1.4074	0.8892	1.5953	0.9080								
0.2842	1.4277	0.8407	1.6202	0.8282								
0.0793	2 4 2 4 9	4 2087	2 2489	0.2311								

Table 5(a). The Avg. and MSE, of different weighted schemes at  $(\delta, \lambda) = (1.5, 2.5)$  and (n, m) =

DWCRTE

Avg.

1.3067

1.0486

	c		0.00/0	0.1057	1 40 46	0.7701	1 2002	0.5410
	<b>S</b> <sub>1</sub>	LLF-I	0.9962	0.1856	1.4846	0.6601	1.7007	0.5410
		LLF-2	1.0976	0.1101	1.6882	0.4660	1.8/38	0.3209
		PRLF	1.1056	0.1024	1.7018	0.4431	1.8873	0.2983
		MLE	1.3135	0.0780	2.3729	3.1072	2.2422	0.2274
		SLF	0.8639	0.4090	1.2871	1.0673	1.4748	1.1918
	\$ <sub>2</sub>	LLF-1	0.7868	0.5359	1.1649	1.2812	1.3432	1.5618
		LLF-2	0.9345	0.3116	1.4074	0.8892	1.5953	0.9080
(0515)		PRLF	0.9491	0.2842	1.4277	0.8407	1.6202	0.8282
(0.0,1.0)		MLE	1.3174	0.0793	2.4249	4.2087	2.2489	0.2311
		SLF	0.8543	0.4305	1.2749	1.1017	1.4584	1.2547
	<b>S</b> <sub>3</sub>	LLF-1	0.7775	0.5607	1.1538	1.3151	1.3272	1.6339
		LLF-2	0.9247	0.3304	1.3942	0.9238	1.5786	0.9628
		PRLF	0.9402	0.3007	1.4157	0.8724	1.6050	0.8763
		MLE	1.3107	0.1344	2.5900	9.0307	2.2375	0.3917
		SLF	0.5819	1.4653	0.9524	1.9576	0.9934	4.2703
	<i>S</i> <sub>4</sub>	LLF-1	0.4553	2.0057	0.8024	2.3463	0.7773	5.8452
		LLF-2	0.6946	1.0729	1.0997	1.6268	1.1857	3.1266
		PRLF	0.7249	0.9285	1.1309	1.5257	1.2374	2.7059
		MLE	0.4189	0.0078	0.7511	0.5409	0.9721	0.0420
		SLF	0.3141	0.0171	0.4534	0.0597	0.7288	0.0921
	<i>S</i> <sub>1</sub>	LLF-1	0.2966	0.0213	0.4210	0.0698	0.6883	0.1145
		LLF-2	0.3310	0.0137	0.4864	0.0517	0.7681	0.0738
		PRLF	0.3328	0.0130	0.4889	0.0497	0.7722	0.0702
		MLE	0.4250	0.0121	0.9170	1.9017	0.9862	0.0654
		SLF	0.2601	0.0374	0.3672	0.1030	0.6036	0.2012
	<i>S</i> <sub>2</sub>	LLF-1	0.2378	0.0459	0.3312	0.1202	0.5518	0.2473
		LLF-2	0.2816	0.0302	0.4039	0.0883	0.6534	0.1628
		PRLF	0.2847	0.0286	0.4079	0.0849	0.6607	0.1539
(1.5,2.5)		MLE	0.4259	0.0120	0.9255	2.0453	0.9882	0.0648
		SLF	0.2591	0.0379	0.3661	0.1043	0.6013	0.2041
	<i>S</i> <sub>3</sub>	LLF-1	0.2371	0.0464	0.3305	0.1212	0.5502	0.2496
		LLF-2	0.2803	0.0308	0.4022	0.0899	0.6503	0.1661
		PRLF	0.2836	0.0291	0.4065	0.0863	0.6581	0.1568
		MLE	0.4289	0.0179	1.1547	4.2872	0.9952	0.0965
		SLF	0.1958	0.0752	0.2797	0.1632	0.4543	0.4047
	<i>S</i> <sub>4</sub>	LLF-1	0.1666	0.0921	0.2391	0.1898	0.3865	0.4958
	-	LLF-2	0.2237	0.0610	0.3211	0.1398	0.5190	0.3283
		PRLF	0.2291	0.0569	0.3272	0.1338	0.5316	0.3066
		•	•	•			•	

(4 -)	C el	Fation - t -	DWCRTE		DWC	DWCRRE		HCE
( <i>t</i> , <i>c</i> )	( <i>t</i> , <i>c</i> ) Sch.	Estimate	Avg.	MSE	Avg.	MSE	Avg.	MSE
		MLE	1.3110	0.0342	2.1955	0.3041	2.2381	0.0995
		SLF	1.1581	0.0580	1.7958	0.2938	1.9770	0.1692
	<i>S</i> <sub>1</sub>	LLF-1	1.1276	0.0706	1.7270	0.3326	1.9250	0.2057
		LLF-2	1.1872	0.0482	1.8640	0.2651	2.0267	0.1405
		PRLF	1.1907	0.0461	1.8710	0.2569	2.0327	0.1344
		MLE	1.3152	0.0380	2.2154	0.3439	2.2453	0.1108
		SLF	1.1294	0.0749	1.7371	0.3508	1.9281	0.2183
	<b>S</b> <sub>2</sub>	LLF-1	1.0968	0.0911	1.6660	0.3996	1.8724	0.2656
		LLF-2	1.1605	0.0619	1.8075	0.3128	1.9811	0.1805
(0 = 1 = 1)		PRLF	1.1645	0.0592	1.8150	0.3028	1.9879	0.1724
(0.5,1.5)		MLE	1.3155	0.0391	2.2181	0.3516	2.2457	0.1140
		SLF	1.1232	0.0810	1.7264	0.3695	1.9174	0.2359
	<b>S</b> <sub>3</sub>	LLF-1	1.0909	0.0979	1.6567	0.4192	1.8623	0.2853
		LLF-2	1.1539	0.0673	1.7955	0.3305	1.9699	0.1961
		PRLF	1.1581	0.0643	1.8034	0.3199	1.9770	0.1873
		MLE	1.3186	0.0432	2.2352	0.3925	2.2510	0.1258
		SLF	1.0939	0.1014	1.6684	0.4316	1.8674	0.2954
	<b>S</b> <sub>4</sub>	LLF-1	1.0580	0.1235	1.5936	0.4944	1.8061	0.3600
		LLF-2	1.1280	0.0834	1.7424	0.3812	1.9256	0.2431
		PRLF	1.1326	0.0795	1.7508	0.3684	1.9335	0.2317
		MLE	0.4198	0.0054	0.6888	0.0451	0.9740	0.0292
	<i>S</i> <sub>1</sub>	SLF	0.3519	0.0082	0.5220	0.0349	0.8166	0.0439
		LLF-1	0.3406	0.0097	0.4983	0.0389	0.7903	0.0522
		LLF-2	0.3630	0.0069	0.5460	0.0321	0.8424	0.0373
		PRLF	0.3637	0.0067	0.5469	0.0312	0.8439	0.0361
		MLE	0.4219	0.0062	0.7046	0.0948	0.9791	0.0335
		SLF	0.3421	0.0102	0.5037	0.0411	0.7937	0.0551
	<i>S</i> <sub>2</sub>	LLF-1	0.3301	0.0121	0.4795	0.0460	0.7660	0.0653
		LLF-2	0.3537	0.0087	0.5282	0.0375	0.8207	0.0467
(1525)		PRLF	0.3545	0.0084	0.5293	0.0365	0.8225	0.0452
(1.3,2.3)		MLE	0.4217	0.0063	0.7094	0.1337	0.9784	0.0341
		SLF	0.3406	0.0107	0.5014	0.0425	0.7903	0.0577
	<b>S</b> <sub>3</sub>	LLF-1	0.3289	0.0126	0.4778	0.0474	0.7632	0.0681
		LLF-2	0.3520	0.0091	0.5252	0.0388	0.8167	0.0491
		PRLF	0.3528	0.0088	0.5265	0.0378	0.8187	0.0476
		MLE	0.4239	0.0073	0.7324	0.2407	0.9837	0.0390
		SLF	0.3301	0.0132	0.4824	0.0493	0.7660	0.0709
	<b>S</b> <sub>4</sub>	LLF-1	0.3173	0.0156	0.4573	0.0553	0.7363	0.0841
		LLF-2	0.3426	0.0111	0.5078	0.0445	0.7949	0.0600
		PRLF	0.3435	0.0108	0.5091	0.0434	0.7971	0.0580

**Table 5(b)**. The Avg. and MSE of different weighted entropy estimates for BXIID under PT-IIC schemes at  $(\delta, \lambda) = (1.5, 2.5)$  and (n, m) = (40, 30)

(( )	6.1		DWG	DWCRTE		CRRE	DWCRHCE	
(t, c)	Sch.	Estimate	Avg.	MSE	Avg.	MSE	Avg.	MSE
		MLE	1.3122	0.0258	2.1817	0.2285	2.2401	0.0751
		SLF	1.2038	0.0334	1.8889	0.1946	2.0550	0.0974
	<i>S</i> <sub>1</sub>	LLF-1	1.1833	0.0385	1.8390	0.2104	2.0199	0.1121
		LLF-2	1.2237	0.0294	1.9385	0.1839	2.0891	0.0858
		PRLF	1.2259	0.0286	1.9432	0.1800	2.0928	0.0832
		MLE	1.3162	0.0300	2.2017	0.2704	2.2469	0.0874
		SLF	1.1725	0.0464	1.8201	0.2450	2.0016	0.1353
	<i>S</i> <sub>2</sub>	LLF-1	1.1502	0.0538	1.7681	0.2688	1.9635	0.1569
		LLF-2	1.1941	0.0404	1.8718	0.2268	2.0384	0.1176
		PRLF	1.1966	0.0391	1.8771	0.2215	2.0427	0.1138
(0.5,1.5)		MLE	1.3166	0.0308	2.2042	0.2768	2.2476	0.0896
		SLF	1.1687	0.0494	1.8136	0.2570	1.9951	0.1440
	<b>S</b> <sub>3</sub>	LLF-1	1.1469	0.0570	1.7631	0.2808	1.9580	0.1660
		LLF-2	1.1898	0.0432	1.8638	0.2387	2.0311	0.1259
		PRLF	1.1925	0.0418	1.8695	0.2331	2.0358	0.1218
		MLE	1.3190	0.0358	2.2210	0.3236	2.2516	0.1042
		SLF	1.1347	0.0678	1.7424	0.3206	1.9370	0.1977
	<i>S</i> <sub>4</sub>	LLF-1	1.1093	0.0792	1.6861	0.3558	1.8937	0.2308
		LLF-2	1.1591	0.0584	1.7983	0.2923	1.9787	0.1701
		PRLF	1.1622	0.0563	1.8045	0.2848	1.9841	0.1641
		MLE	0.4174	0.0039	0.6745	0.0313	0.9686	0.0213
	<i>S</i> <sub>1</sub>	SLF	0.3685	0.0048	0.5522	0.0233	0.8551	0.0261
		LLF-1	0.3605	0.0055	0.5345	0.0251	0.8365	0.0298
		LLF-2	0.3764	0.0043	0.5701	0.0221	0.8733	0.0231
		PRLF	0.3767	0.0042	0.5706	0.0217	0.8742	0.0226
		MLE	0.4195	0.0048	0.6845	0.0393	0.9734	0.0258
		SLF	0.3573	0.0066	0.5304	0.0294	0.8291	0.0358
	<b>S</b> <sub>2</sub>	LLF-1	0.3488	0.0076	0.5122	0.0320	0.8094	0.0409
		LLF-2	0.3657	0.0058	0.5488	0.0275	0.8485	0.0315
(1 5 2 5)		PRLF	0.3662	0.0057	0.5495	0.0270	0.8496	0.0307
(1.3,2.3)		MLE	0.4192	0.0048	0.6839	0.0393	0.9728	0.0259
		SLF	0.3568	0.0069	0.5297	0.0302	0.8279	0.0369
	<b>S</b> <sub>3</sub>	LLF-1	0.3486	0.0078	0.5122	0.0327	0.8089	0.0420
		LLF-2	0.3648	0.0061	0.5474	0.0284	0.8465	0.0327
		PRLF	0.3654	0.0059	0.5483	0.0278	0.8478	0.0319
		MLE	0.4211	0.0058	0.6993	0.0829	0.9772	0.0313
		SLF	0.3442	0.0093	0.5060	0.0376	0.7988	0.0499
	<i>S</i> <sub>4</sub>	LLF-1	0.3349	0.0106	0.4866	0.0411	0.7770	0.0573
		LLF-2	0.3534	0.0081	0.5255	0.0347	0.8201	0.0437
		PRLF	0.3541	0.0079	0.5264	0.0340	0.8216	0.0426

**Table 5(c).** The Avg. and MSE, of different weighted entropy estimates for BXIID under PT-IIC schemes at  $(\delta, \lambda) = (2.5, 1.5)$  and (n, m) = (60, 40)

( <i>t</i> , <i>c</i> )	Sch.	Estimate	DWCRTE		DWCRRE		DWCRHCE	
			Avg.	MSE	Avg.	MSE	Avg.	MSE
(0.5,1.5)	<i>S</i> <sub>1</sub>	MLE	1.3126	0.0213	2.1735	0.1882	2.2407	0.0620
		SLF	1.2278	0.0238	1.9409	0.1514	2.0960	0.0694
		LLF-1	1.2117	0.0265	1.9003	0.1590	2.0685	0.0773
		LLF-2	1.2435	0.0218	1.9812	0.1470	2.1228	0.0635
		PRLF	1.2450	0.0213	1.9846	0.1448	2.1254	0.0621
	<b>S</b> 2	MLE	1.3142	0.0224	2.1803	0.1996	2.2434	0.0653
		SLF	1.2190	0.0263	1.9201	0.1614	2.0809	0.0766
		LLF-1	1.2026	0.0294	1.8794	0.1707	2.0530	0.0856
		LLF-2	1.2349	0.0239	1.9606	0.1555	2.1081	0.0696
		PRLF	1.2365	0.0233	1.9641	0.1530	2.1108	0.0680
	<b>S</b> 3	MLE	1.3145	0.0229	2.1820	0.2035	2.2440	0.0666
		SLF	1.2161	0.0276	1.9141	0.1672	2.0760	0.0803
		LLF-1	1.1999	0.0308	1.8740	0.1769	2.0484	0.0897
		LLF-2	1.2319	0.0250	1.9540	0.1608	2.1029	0.0729
		PRLF	1.2335	0.0245	1.9577	0.1582	2.1057	0.0713
	\$ <sub>4</sub>	MLE	1.3156	0.0238	2.1871	0.2132	2.2458	0.0695
		SLF	1.2088	0.0297	1.8970	0.1757	2.0635	0.0866
		LLF-1	1.1919	0.0334	1.8555	0.1872	2.0347	0.0973
		LLF-2	1.2252	0.0268	1.9383	0.1678	2.0916	0.0781
		PRLF	1.2269	0.0262	1.9419	0.1649	2.0944	0.0762
(1.5,2.5)	<i>S</i> <sub>1</sub>	MLE	0.4170	0.0032	0.6694	0.0252	0.9675	0.0174
		SLF	0.3775	0.0035	0.5698	0.0181	0.8760	0.0188
		LLF-1	0.3711	0.0039	0.5551	0.0191	0.8611	0.0210
		LLF-2	0.3839	0.0032	0.5847	0.0176	0.8908	0.0172
		PRLF	0.3841	0.0031	0.5850	0.0173	0.8913	0.0169
	<b>S</b> <sub>2</sub>	MLE	0.4177	0.0035	0.6726	0.0274	0.9693	0.0187
		SLF	0.3743	0.0039	0.5632	0.0196	0.8685	0.0209
		LLF-1	0.3678	0.0043	0.5485	0.0207	0.8534	0.0233
		LLF-2	0.3807	0.0035	0.5781	0.0189	0.8835	0.0190
		PRLF	0.3810	0.0035	0.5784	0.0186	0.8840	0.0187
	<b>S</b> <sub>3</sub>	MLE	0.4177	0.0035	0.6727	0.0278	0.9692	0.0190
		SLF	0.3735	0.0040	0.5617	0.0201	0.8667	0.0217
		LLF-1	0.3671	0.0045	0.5473	0.0213	0.8518	0.0242
		LLF-2	0.3798	0.0037	0.5763	0.0194	0.8813	0.0197
		PRLF	0.3801	0.0036	0.5767	0.0191	0.8820	0.0194
	<i>S</i> <sub>4</sub>	MLE	0.4184	0.0038	0.6759	0.0300	0.9709	0.0202
		SLF	0.3705	0.0044	0.5557	0.0215	0.8598	0.0237
		LLF-1	0.3639	0.0049	0.5407	0.0228	0.8443	0.0265
		LLF-2	0.3771	0.0040	0.5708	0.0206	0.8751	0.0215
		PRLF	0.3774	0.0039	0.5712	0.0203	0.8757	0.0211

**Table 5(d).** The Avg. and MSE of different weighted entropy estimates for BXIID under PT-IIC schemes at( $\delta$ ,  $\lambda$ ) = (2.5,1.5) and (n, m) = (60, 50)

It is evident from the tabulated result values that:

- 1- As *n* and *m* increase, the MSE of the Bayes estimate gradually decreases.
- 2- In comparison to other estimates, the MSE of all estimates based on DWCRTE frequently yields the smallest values.
- 3- As can be seen in Tables 4 (a) to 4 (d), in the majority of the cases, the precision measures of DWCRTE, DWCRRE and DWCRHCE under PRLF and LLF-2 are preferable to the corresponding estimates under LLF-1 and SLF for all schemes.
- 4- At true values  $T^*(c) = 0.3434$  and 1.3621, the BEs of  $T^*(c)$  under all loss functions are preferred over the other entropy measures for all schemes (see Tables 4(a) and 4(d)).
- 5- The MLEs and BEs of T\*(c), R\*(c) and H\*(c) under different loss functions are decreasing as n and m increase from (40, 20) to (60, 50) (see Tables 4(a) and 4(d)).
- 6- The MSE of all BEs of  $T^*(c)$  gets the smallest values compared to the others for all schemes at (n, m) = (60, 40) and (t, c) = (0.5, 1.5) and (1.5, 2.5) (see Table 5(c)).
- 7- In most cases, the MSEs of all estimates have the largest values in the case of  $S_3$  and  $S_4$  compared to other schemes, at (n, m) = (60, 50), (t, c) = (0.5, 1.5) and (1.5, 2.5) (see Table 5(d)).
- 8- The MSE of  $T^*(c)$  gets the smallest values under all loss functions in the case of  $S_1$  at m = 20, n = 40 (see Figure 4).
- 9- At true values of  $H^*(c) = 0.7968$  and  $T^*(c) = 0.3434$ , it can be observed that the BE of DWCRHCE gets the largest MSE, while the BE of DWCRTE gets the smallest value for all loss functions in Sch.1 (see Figure 5).









Source: created by researchers utilizing Microsoft Excel.

- 10-The MSE of the BE for  $R^*(c)$  based on S<sub>1</sub> has the biggest values at m = 50, n = 60 and t = 0.5 (see Figure 6).
- 11-The MSE of the DWCRTE estimate under different loss functions based on  $S_1$  typically produces the smallest values when compared to other estimates at m = 50, n = 60 and t=1.5 (see Figure 7).





**Figure 6.** MSE of different Entropy estimates at t = 0.5, m = 50, and n = 60 t = 1.5Source: created by researchers utilizing Microsoft Excel.

- 12-It can be observed from Figure 8 that the MSE of  $\hat{T}^*(c)$  under PRLF takes the smallest values compared to the others in Sch. 4 at m = 20, n = 40, and t = 0.5.
- 13-At m = 20, n = 40, where the true value of  $\hat{T}^*(c) = 0.3434$ , the MSE of  $\hat{T}^*(c)$  under LLF-2 takes the smallest values compared to the others under S<sub>4</sub> (Figure 9).
- 14-When compared to the other estimates from  $S_2$ ,  $S_3$ , and  $S_4$ , the MSE of all entropy estimates based on  $S_1$  often has the smallest values. The majority of entropy estimates (MLE, SLF, LLF-1, LLF-2, and PRLF) show a slight decrease as *t* increases.
- 15-It should be highlighted that, in comparison to the other estimates based on *S*<sub>4</sub>, *S*<sub>3</sub> often produces the shortest MSE outcomes.



0,2 0,15 0,05 0 SLF LLF-1 LLF-2 PRLF



**Figure 9**. MSE of DWCRTE for different loss functions when *t* =1.5

Source: created by researchers utilizing Microsoft Excel.



16-As can be seen in Figures 10 and 11, as *n* and *m* increase, the MSEs of  $R^*(c)$  under different loss functions are decreasing.



**Figure 10**. MSE of DWCRRE for different estimates under  $S_3$  at  $\delta = 1.5$ ,  $\lambda = 2.5$  and t = 1.5



Source: created by researchers utilizing Microsoft Excel.

# 8. Concluding Remarks

This article introduces the DWCRHC as an additional measure of uncertainty surrounding the residual lifetime function, particularly relevant in fields like survival analysis and reliability. The DWCRHC measure is formally defined within this work. We investigate the estimation of the DWCRHC, along with its related measures, DWCRRE and DWCRTE, for the BXIID under PT-IIC. Maximum and Bayesian estimation methods are employed. For the Bayesian estimation, we utilize the MCMC approach with the M-H algorithm, assuming a gamma prior distribution and considering three different loss functions. The article features the application, simulation studies, and an evaluation of the accuracy of the DWCRTE, DWCRRE, and DWCRHCE estimates for the BXIID.

Simulation results demonstrate that the BE of DWCRTE converges to the true value as the sample size increases. Generally, BEs under the PRLF exhibit the lowest MSE values, followed by the LLF-2, making them preferable over competing estimates. Furthermore, Sch. 1, compared to other schemes, often yields the lowest MSE values, followed in most cases by Sch. 2. The conclusions drawn from the simulated data are corroborated by the examination of actual data, particularly the water capacity data from the Shasta reservoir. These results are helpful in making well-informed decisions about the management of water resources. Future research could explore the application of the E-Bayesian technique to estimate other uncertainty metrics, such as dynamic weighted cumulative residual Shannon entropy.

## References

- Abd-Elfattah, A. M., Hassan, A. S. and Nassr, S. G., (2008). Estimation in step-stress partially accelerated life tests for the Burr Type XII distribution using type I censoring. *Statistical Methodology*, 5(6), pp. 502–514.
- Abo-Eleneen, Z. A., (2011). The entropy of progressively censored samples. *Entropy*, 13 (2), pp. 437–49.
- Ahmadini, A. H., Hassan, A. S., Zaky, A. N. and Alshqaq, S. S., (2020). Bayesian inference of dynamic cumulative residual entropy from Pareto II distribution with Application to COVID-19. AIMS Mathematics, 6(3), pp. 2196–2216.
- Al-Babtain, A. A., Hassan, A.S., Zaky, A. N., Elbatal, I. and Elgarhy, M., (2021). Dynamic cumulative residual Rényi entropy for Lomax distribution: Bayesian and non-Bayesian methods. *AIMS Mathematics*, 6(3), pp. 3889–3914.
- AlmarashiI, M., Algarni, A., Hassan, A. S., Zaky, A. S. and Elgarhy, M., (2021). Bayesian analysis of dynamic cumulative residual entropy for Lindley distribution. *Entropy*, 23, 1256. https://doi.org/10.3390/ e23101256.
- Alyami, S. A., Hassan, A. S., Elbatal, I., Elgarhy, M., A. R. and El-Saeed, A. R., (2023). Bayesian and non- Bayesian estimates of the DCRTE for moment exponential distribution under progressive censoring type II. Open Physics. 21, 20220264.
- Asadi, M., Zohrevand, Y., (2007). On the dynamic cumulative residual entropy. *Journal* of *Statistical Planning and Inference*, 137, pp. 1931–1941.
- Balakrishnan, N., Aggrawala, R., (2000). *Progressive Censoring, Theory, Methods and Applications*. Birkhauser, Boston.
- Baratpour, S., Ahmadi, J. and Arghami, N. R., (2007). Entropy properties of record statistics. *Statistical Papers*, 48, pp. 197–213.
- Belis, M., Guiasu, S., (1968). A quantitative-qualitative measure of information in cybernetic systems. *IEEE Transactions on Information Theory*, IT-4, pp. 593– 594.
- Burr, W. I., (1942). Cumulative frequency functions. Annals of Mathematical Statistics, 13(2), pp. 215–232.
- Belzunce, F., Navarro, J., Runiz, J. M. and Aguila, Y., (2004). Some results on residual entropy function. *Metrika*, 59, pp. 147–161.
- Cho, Y., Sun, H. and Lee, K., (2015). Estimating the entropy of a Weibull distribution under generalized progressive hybrid censoring. *Entropy*, 17, pp. 102–122.

- Cohen, A. C., (1963). Progressively censored samples in life testing. Technometrics, 5(3), pp. 327–339.
- Di Crescenzo, A. D, Longobardi, M., (2006). On weighted residual and past entropies, Scientiae Mathematicae Japonicae, 64(3), pp. 255–266.
- Evans, R. A., Simons, G., (1975). Research on statistical procedures in reliability engineering. ARL TR 75-0154, AD A013687.
- Guiasu, S., (1986). Grouping data by using the weighted entropy. *Journal of Statistical Planning and Inference*, 15, pp. 63–69.
- Gupta, P. L., Gupta, R. C. and Lvin, S. J., (1996). Analysis of failure time data by Burr distribution. *Communications in Statistics-Theory and Methods*, 25, pp. 2013–2024.
- Hassan, A. S., Zaky, A. N., (2021). Entropy Bayesian estimation for Lomax distribution based on record. *Thailand Statistician*,19(1), pp. 96–115.
- Hassan, A. S., Zaky, A. N., (2019). Estimation of entropy for inverse Weibull distribution under multiple censored data. *Journal of Taibah University for Science*, 13, pp. 331–337.
- Hassan, A. S., Assar, S. M. and Ali, K.A., (2024a). Efficient estimation of the Burr XII distribution in presence of progressive censored samples with Binomial random removal. *Thailand Statistician*, 22(1), pp. 121–141.
- Hassan, A. S., Alsadat, N., Balogu, O. S. and Helmy, B. A., (2024b). Bayesian and non-Bayesian estimation of some entropy measures for a Weibull distribution. *AIMS Mathematics*, 9(11), 32646–32673. DOI: 10.3934/math.20241563
- Hassan, A. S., Elsherpieny, E. A. and Mohamed, R. E., (2022). Estimation of information measures for power-function distribution in presence of outliers and their applications. *Journal of Information and Communication Technology*, 21 (1), pp. 1–25.
- Havrda, J., Charvat, F., (1967). Quantification method of classification process: Concept of structural α-entropy, *Kybernetika*, 3, pp. 30–35.
- Helmy, B.A., Hassan, A. S. and El-Kholy, A. K., (2021). Analysis of uncertainty measure using unified hybrid censored data with applications. *Journal of Taibah University for Science*, 15(1), pp. 1130–1143.
- Helmy, B. A., Hassan, A. S. and El-Kholy, A. K., (2023). Analysis of Uncertainty Weighted Measures for Pareto II distribution. *Reliability Theory & Applications*, 18(3), pp. 81–196.

- Kayal, S., Balakrishnan, N., (2023). Weighted fractional generalized cumulative past entropy and its properties. *Methodology Computing Applied Probability*, 25, p 61. https://doi.org/10.1007/s11009-023-100350
- Khammar, A.H., Jahanshahi, A. M. A., (2018). On weighted cumulative residual Tsallis entropy and its dynamic version. *Physica A: Statistical Mechanics*, 491, pp. 678– 692.
- Lee, K., (2017). Estimation of entropy of the inverse Weibull distribution under generalized progressive hybrid censored data. *Journal of the Korean Data & Information Science Society*, 28(3), pp. 659–668.
- Li, X., Shi, Y., Wei, J. and Chai, J., (2007). Empirical Bayes estimators of reliability performances using LINEX loss under progressively type-II censored samples. *Mathematics and Computers in Simulation*, 73(5), pp. 320–326.
- Misagh, F., Yari, G. H., (2011). On weighted interval entropy. *Statistics & Probability Letters*, 81, pp. 188–194.
- Mohamed, M. S., (2022). On cumulative residual Tsallis entropy and its dynamic version of concomitants of generalized order statistics. *Communications in Statistics-Theory and Methods*, 51(8), pp. 2534–2551.
- Mousa, M. A., Jaheen, Z. F., (2002). Statistical inference for the Burr model base on progressively censored data. *Computers & Mathematics with Applications*, 43, pp. 1441–1449.
- Nair, R. S., Sathar and E. I. A., (2024). Bivariate dynamic weighted cumulative residual entropy. *Japanese Journal of Statistics and Data Science*, 7, pp. 83–100 (2024). https://doi.org/10.1007/s42081-023-00232-z
- Nanda, A. K., Paul, P., (2006). Some results on generalized residual entropy, Information Sciences, 176, pp. 27–47.
- Nadar, M., Papadopoulos and A. Kızılaslan, F., (2013). Statistical analysis for Kumaraswamy's distribution based on record data. *Statistical Papers*, 54, pp. 355– 369.
- Navarro, J., Del Aguila, Y. and Ruiz, J. M., (2001). Characterizations through reliability measures from weighted distributions. *Statistical Papers*, 42, pp. 395–402.
- Panahi, H., Sayyareh, A., (2014). Parameter estimation and prediction of order statistics for the Burr type XII distribution with type II censoring, *Journal of Applied Statistics*, 41, pp. 215–232.

- Qin, X., Gui, W., (2020). Statistical inference of Burr-XII distribution under progressive type-II censored competing risks data with binomial removals. *Journal of Computational and Applied Mathematics*, 378, 112922.
- Rao, M., Chen, Y., Vemuri, B. C. and Wang, F., (2004). Cumulative residual entropy: a new measure of information. *IEEE Transactions Information Theory*, 50, pp. 1220– 1228.
- Renjini, K. R., Abdul Sathar, E. I. and Rajesh, G., (2016a). Bayes estimation of dynamic cumulative residual entropy for Pareto distribution under type-II right censored data. *Applied Mathematical Modelling*, 40, pp. 8424–8434.
- Renjini, K. R., Abdul Sathar, E. I. and Rajesh, G., (2016b). A study of the effect of loss functions on the Bayes estimates of dynamic cumulative residual entropy for Pareto distribution under upper record values. *Journal of Statistical Computation and Simulation*, 86, pp. 324–339.
- Renjini, K. R., Abdul Sathar, E. I. and Rajesh, G., (2018). Bayesian estimation of dynamic cumulative residual entropy for classical Pareto distribution. *American Journal of Mathematical and Management Sciences*, 37, pp. 1–13.
- Rény, A., (1961). On measures of entropy and information in proceeding of the fourth Berkeley Symposium on Mathematical Statistics and Probability, University of California Press: Berkeley, CA, USA, 1, pp. 547–561.
- Sati, M. M., Gupta, N., (2015). Some characterization results on dynamic cumulative residual Tsallis entropy, *Journal of Probability and Statistics*. 23 http://dx.doi.org/ 10.1155/2015/694203.ArticleID 694203, p. 8.
- Shannon, C. E., (1948). A mathematical theory of communication. *Bell System Technical Journal*, 27, pp. 379–423.
- Smitha, S., Sudheesh, K. K. and Sreedevi, E. P., (2024). Dynamic cumulative residual entropy generating function and its properties. *Communications in Statistics -Theory and Methods*, 53(16), pp. 5890–5909.
- Soliman, A. A., (2005). Estimation of parameters of life from progressively censored data using Burr-XII model. *IEEE Transactions on Reliability*, 54(1), pp. 34–42.
- Sunoj, S.M., Linu, M. N., (2012). Dynamic cumulative residual Rényi's entropy. Statistics, 46(1), pp. 41–56.
- Sunoj, S.M., Maya, S. S., (2006). Some properties of weighted distribution in the context of repairable systems. *Communications in Statistics—Theory and Methods*, 35(2), pp. 223–228.

- Tsallis, C., (1988). Possible generalization of Boltzmann-Gibbs statistics. *Journal* of *Statistical Physics*, 52, pp. 479– 487.
- Wang, M-C., (1996). Hazards regression analysis for length-biased data. *Biometrika*, 83, pp. 343-354.
- Wang, X., Gui, W., (2021). Bayesian estimation of entropy for Burr Type XII distribution under progressive Type-II censored data. *Mathematics*, 9, p. 313. https://doi.org/10.3390/math9040313.
- Wingo, D. R., (1993). Maximum likelihood methods for fitting the Burr type XII distribution to multiply (progressively) censored life test data. *Metrika*, 40, pp. 203–210.
- Wu, J. W., Yu, H. Y., (2005). Statistical inference about the shape parameter of the Burr type XII distribution under the failure-censored sampling plan. *Applied Mathematics and Computation*, 163(1), pp. 443–482.